

REVIEW

Open Access



Factors influencing performance and injury risk in elite female Gaelic team sport players and future research directions: a narrative review

John David Duggan^{1*}, Kieran Collins² and Karen Keane¹

Abstract

Background: Sports science research in elite female Gaelic team sports has increased in recent years, but still a large disparity exists between the volume of studies involving male and female players. As a consequence of this, it is difficult for practitioners to develop an evidence-based approach when working with female players.

Main body: In this review, we discuss the current research available in elite female Gaelic team sports with focus on seven specific areas including physical and physiological demands, anthropometric and performance characteristics, injury risk, nutritional considerations, and female physiology. There appears to be unique physical demands data in match play across positions in Camogie, however, there is currently no comparative data available in ladies Gaelic football. Similarly, there is no research available on the physiological demands of both elite female Gaelic team sports. According to existing literature, performance characteristics such as speed and power are lower in this population compared to other elite female team sports. Although data is limited, the anthropometric characteristics of elite female Gaelic team sport players appear homogenous with some positional differences observed at a sub-elite level. Previous research has demonstrated a high prevalence of lower limb injuries in female elite Gaelic team sports and the provision of quality, evidence-based strength & conditioning could help mitigate these injury risks. Female Gaelic team sport players have been shown to have poor nutrition knowledge and inadequate intakes of micronutrients. Finally, although menstrual cycle phase and oral contraceptives have been shown to influence performance in other female intermittent sports, to date there has not been any research carried out in elite female Gaelic team sport players.

Conclusions: It is evident that limited research has been carried out on elite female Gaelic sport players. More up-to-date, high-quality investigations are needed to address the research gaps, which in turn should enable practitioners in the field to apply sound, evidence-based practice/theory when working with this population.

Keywords: Elite female Gaelic sports, Gaelic games, Performance, Nutrition, Injury, Anthropometrics

Background/introduction

Camogie [kuh mow gee] and ladies Gaelic football are two of the most popular female participation-based sports in Ireland. Furthermore, as the Irish diaspora settle around the world and the increasing globalization of sport, both games are now played in every corner of the globe. Despite the amateur ethos of Gaelic games,

*Correspondence: john.duggan@atu.ie

¹ Department of Sport, Exercise & Nutrition, School of Science & Computing, Atlantic Technological University Galway, Dublin Road, Galway, Ireland
Full list of author information is available at the end of the article



© The Author(s) 2022. **Open Access** This article is licensed under a Creative Commons Attribution 4.0 International License, which permits use, sharing, adaptation, distribution and reproduction in any medium or format, as long as you give appropriate credit to the original author(s) and the source, provide a link to the Creative Commons licence, and indicate if changes were made. The images or other third party material in this article are included in the article's Creative Commons licence, unless indicated otherwise in a credit line to the material. If material is not included in the article's Creative Commons licence and your intended use is not permitted by statutory regulation or exceeds the permitted use, you will need to obtain permission directly from the copyright holder. To view a copy of this licence, visit <http://creativecommons.org/licenses/by/4.0/>. The Creative Commons Public Domain Dedication waiver (<http://creativecommons.org/publicdomain/zero/1.0/>) applies to the data made available in this article, unless otherwise stated in a credit line to the data.

inter-county (elite) female Gaelic team sport players complete up to six tactical, technical and physical training sessions per week, consisting of combination of both pitch and resistance training-based sessions [1, 2]. During the competitive season, elite inter-county female Gaelic team sport players compete in two major competitions, the National League, and the All-Ireland Championship. The National League runs from January to April, whilst the All-Ireland Championship runs from May to July each year. From a competition/training perspective, during the peak competitive phases of the National League and All-Ireland, female athletes may partake in games on a weekly/fortnightly basis depending on progress through each of the major competitions.

In both female Gaelic team sports, the game model of both sports is similar [3]. A team consists of 15 players, with the possibility of using five substitutes [4]. Each team includes a goalkeeper, two lines of three defensive players (full-back and half-back), two midfielders, and two lines of attacking players (half forward and full forward). Games at elite inter-county level comprise of two 30-min halves played on a rectangular pitch, 145 m in length and 90 m in width [5]. The principles of play of both female Gaelic team sports are to disrupt and unbalance the defense by striking the Gaelic football or the solid leather *slíotar* [slit er] (camogie) through the oppositions goalposts (similar to rugby goalposts), either below the cross bar for a goal (three points) or above for a point [1]. In ladies Gaelic football (similar principles of play to Australian Rules Football and basketball), the ball (diameter 680 mm, mass 480 g) is spherical, and heavier than the ball used in soccer. The skills of the ladies Gaelic football include high catching/fielding, handling the ball, kicking the football over a range of distances, solo running with the ball, passing the ball by hand, and blocking and intercepting [4]. Successful ladies Gaelic football teams were observed to have significantly superior ability to gain and use possession [6]. In camogie, (similar principles of play to field hockey & Lacrosse), an ash stick (hurley) is used to strike the *slíotar* (diameter 69–72 mm, mass 110–120 g) over variable distances. The skills of camogie include high catching/fielding, aerial duels, striking the *slíotar* over short and long distances, passing the *slíotar* by hand, solo running with the *slíotar* balanced on the hurley, blocking opponents using the hurley and intercepting the *slíotar* with the hurley or by hand [1, 4].

There has been an increase in both participation and professionalization of female team sports in the past decade [6, 7]. Despite this, there appears to be an underrepresentation of sports science research focused on female athletes [8, 9]. It has been reported that only 4% of research conducted in sport science and medicine includes female-only cohorts [10, 11]. This disparity

influences practitioners to apply evidence developed in male team athletes' environments to the female team sport domain, which has the potential to lead to erroneous assumptions [9, 12]. Progressively, there has been recent research efforts to compile evidence on the match demands, physiological characteristics and health and performance considerations of female athletes including soccer [13, 14], rugby 7 s [15] and AFL [16]. However, no such source exists for female Gaelic team sports. Therefore, the primary aim of this review is to summarize existing literature relating to elite female Gaelic team sports athletes including physical and physiological demands, anthropometric and performance characteristics, injury risk, nutritional considerations, and female physiology. In doing so, the authors aim to highlight what is currently known on these specific topics and identify gaps in the research. The review will focus on research conducted in elite female and sub elite Gaelic team sports athletes with comparisons made to other female team sports and male Gaelic team sport athletes when appropriate.

Physical demands

The use of microtechnology devices such as global positioning systems (GPS) has become ubiquitous in team sports settings to enable sports practitioners to collect comprehensive and real-time data during training and competition [17]. The GPS data can aid practitioners to objectively quantify the physical demands of competition, examine individual positional workloads and establish training intensities [18]. Despite the growing number of GPS research in the male versions of Gaelic team sports [19–23], research in elite female Gaelic team sports is still at the emergent stage.

Currently, just four peer reviewed studies have been conducted on the physical demands of elite female camogie players. The first study demonstrated that in 60 min of camogie match-play, elite players covered 5881 ± 906 m in total, 546 ± 259 m in high-speed running, and 183 ± 130 m sprinting [24]. A second study reported that players covered 5800 ± 900 m in total distance, 733 ± 245 m in high-speed running and 206 ± 93 m in sprinting [25]. Positional and running temporal demands measured during 5 min time epochs in elite camogie suggested that midfielders had the greater total and sprint distance decrements and defenders also exhibited total and relative decrements in match-play [26]. However, caution is warranted using this approach, as it may underprepare athletes for competition and not be representative of the complex demands of match play [27]. This data is comparable to other female team sports including elite female AFL where players, across the range of positions covered 6.0–7.0 km at $95.0\text{--}126.0$ m min^{-1} during 80 min of match-play [16].

However, the very-high speed running ($>18 \text{ km h}^{-1}$) and sprint distances ($>20 \text{ km h}^{-1}$) were marginally less for these AFL players when compared to camogie, $\sim 417 \text{ m}$ and $\sim 159 \text{ m}$ respectively. Furthermore, it has been shown in camogie that full backs cover less total distance (m) and relative distance (m min^{-1}) compared to other positions, whilst half backs and half forwards had higher sprint distances when compared to other playing positions [24]. This is similar to data in elite female field hockey where forwards were reported to have the highest peak output for relative high-speed distance ($119.3 \pm 19.7 \text{ m min}^{-1}$) in comparison to the defenders ($100.7 \pm 19.7 \text{ m min}^{-1}$) [28]. The maximal velocity reported in elite camogie players has been previously reported as $24.9 \pm 1.6 \text{ km h}^{-1}$ [24], $24.7 \pm 1.26 \text{ km h}^{-1}$ [20] and most recently $25.4 \pm 1.5 \text{ km h}^{-1}$ [29], respectively. These values are slightly higher in comparison to elite Irish female rugby players where maximal velocity has been reported as $23.4 \pm 2.52 \text{ km h}^{-1}$ [30].

Currently, there appears to be no peer-reviewed GPS data for elite female Gaelic football. An early study using time motion analysis investigated player locomotion in elite female Gaelic football and concluded that players spend more time standing still with less time in high intensity activities in comparison to males, with no performance decrements observed across halves of play [31]. However, this method has since been described as outdated. As a result of the paucity of data in ladies Gaelic football, practitioners and coaches rely heavily on data from other female team sports and from the male game, which has received criticism [32]. The use of absolute thresholds is a common example of this, as values from men's Gaelic football have been adopted and applied in the female game [24]. It has previously been suggested that employing male-related speed velocity zones thresholds to the female team sports context could result in an underestimation of workloads [33, 34], and as such female specific high-speed running velocity thresholds have been recommended due to the physiological sex differences in physical fitness/capacity [29, 33–37]. Therefore, an up-to-date analysis of physical demands of elite female Gaelic football using GPS is needed. This information would aid practitioners to design sport and position specific training programs which mimic the unique demands of the game.

Physiological demands

The use of heart rate (HR) measurement has been suggested previously as an indirect measure of exercise intensity and has become a widespread practice in female team sports [13, 38]. However, despite this there has been no research on HR indices in either elite female Gaelic team sports. The application of HR measures during

training and match-play within female Gaelic team sports would ensure that athletes receive an adequate internal training load stimulus and can provide useful feedback in maintaining or improving fitness capacities in preparing for the demands of match-play [32]. HR monitoring could also enable practitioners to design training units to potentially replicate the demands of the game and provide an individualized aerobic stimulus [39, 40]. In elite female soccer, small-sided games elicited a higher HR response ($>85 \text{ HR}_{\text{max}}$) when compared to medium- and large-conditioned games [39]. This information could be used by Gaelic game practitioners to manipulate the constraints within a modified game to elicit a specific physiological adaptation whilst continually developing the technical and tactical elements of the game [41]. It has previously been shown in elite female soccer players the average HR during competitive games ranged between 152 and 186 b min^{-1} , the equivalent ~ 80 and 90% of HR_{peak} [35, 38], with no differences across positions. In contrast, during competitive match play in elite female hockey players, HR_{peak} was reported at $199 \pm 1 \text{ b min}^{-1}$, with an average HR intensity of 86 ± 7.8 of HR_{peak} . Positional differences were also reported with defenders spending more time at $>85\% \text{ HR}_{\text{peak}}$ when compared to forwards ($p \leq 0.001$) [39]. There is a dearth of the research available on the physiological demands in elite female Gaelic team sports. Further research is needed to examine if HR values are similar and/or if positional differences exist in female Gaelic team sports.

A well-developed aerobic system is required for team sport athletes to adequately recover between periods of play and between periods of maximal and submaximal work [42]. Limited data exists on the performance profile of elite female Gaelic team sports athletes. The estimated aerobic power values for successful elite female Gaelic footballers tend to be high, supporting the belief that there is a large aerobic contribution to playing the game. Keane et al. [43] reported a mean value of $49.9 \pm 4.2 \text{ mL kg}^{-1} \text{ min}^{-1}$ for elite female Gaelic footballers. The values compare favorably to elite female soccer ($51.9 \pm 5.1 \text{ mL kg}^{-1} \text{ min}^{-1}$) and female Australian football ($50.4 \pm 6.9 \text{ mL kg}^{-1} \text{ min}^{-1}$) players [13, 44]. Tucker and Reilly [45] reported that the mean of female sub-elite Gaelic football players were slightly lower at $42.0 \pm 6.8 \text{ mL kg}^{-1} \text{ min}^{-1}$. At present, there is no comparative data for elite camogie players.

Performance characteristics

In addition to a well-developed aerobic system, strength, speed and power are all essential bio-motor qualities of female Gaelic team sports [46]. The development of muscular strength and power is an important attribute in elite female Gaelic games athletes to enable them

to tolerate the physical demands of the games [43]. Recently isometric strength, using an isometric mid-thigh pull (IMTP) test reported the average peak force was 1938.46 ± 300.17 N and average relative force as 28.72 ± 3.11 N/kg in elite camogie athletes [47]. This is in comparison to IMTP results in elite female rugby union in which the backs and forwards average peak force as 2560.8 N and 2729.8 N respectively [48]. The average relative force was 36.4 N/kg for the forwards and 31.8 N/kg for backs. This could be due to the superior training age of the elite female rugby union athletes and the unique physical characteristics of the game. There is a myriad of research to suggest the importance of resistance training for female athletes to enhance performance and mitigate injury risks [49–55]. Furthermore, enhanced muscular strength for athletic performance has been advocated by Suhomel et al. [49, 50] who suggests that it correlated with rate of force development (RFD), mechanical power output, and sports specific movements (running, jumping, striking) as well as enhanced ability to perform on-field sports specific skills.

Strength training programs for female Gaelic team sports athletes, which includes weightlifting, ballistic, complex training and plyometric movements enhances neural drive, neural activation rates and inter-muscular coordination [1]. It was demonstrated that females who engaged in a 20-week, multi-joint, strength-based training improved strength (neural adaptations, ($p < 0.05$) with no change in muscle cross sectional area [56]. Advanced training methods such as complex training have been shown to improve athletic capabilities in female athletes with a high training age [57]. From a training specificity perspective, complex training has been demonstrated to improve running economy in division 1 female soccer player [57]. Further research is needed to ascertain the efficacy of specific strength training modalities in elite female Gaelic team sports athletes.

Speed characteristics

Rapid acceleration and sprinting ability are important performance characteristics which allow Gaelic team sports players to reach the ball before the opposition; hence, sprint tests are routinely administered to evaluate performance [58, 59]. In elite camogie, match play sprint and high-speed running distances have been reported as 546 ± 259 m and 183 ± 130 m respectively [24]. O'Grady et al. [29] investigated sprint performance in elite camogie and reported that total sprint distance was 162 ± 102 m with the number of sprints accumulated were 9 ± 5 *n*. The average sprint duration was 3 ± 1 s during competitive, elite match-play. There was also greater number of sprints performed < 20 m in elite, inter-county match-play (7 ± 3 *n*) compared to > 20 m (3 ± 2 *n*) [29]. The mean

length of sprints during elite match-play was reported as 17 ± 4 m. The sprints accumulated between 80%–90% of relative velocity was reported as 6 ± 3 *n* and $> 90\%$ 3 ± 3 *n* during in elite match-play [29]. From a positional perspective, midfielders and half-forwards had the greatest total sprint distance (208 ± 77 & 200 ± 143 m) and peak speed (26.0 ± 1.5 & 25.7 ± 1.9 km h⁻¹) in elite camogie match-play [29]. Tucker and Reilly [45] reported mean 30 m sprint times among elite female Gaelic footballers (5.2 ± 0.2 s). Values for elite camogie players with respect of the shorter distances of 5 (1.20 ± 0.08 s), 10 (2.02 ± 0.08 s) and 20 m (3.45 ± 0.11 s) have recently been reported in the literature [48]. This is in comparison to elite Irish female soccer players where sprint times were reported as 10 m (1.89 ± 0.08 s), 20 m (3.26 ± 0.17 s) and 30 m (4.57 ± 0.26) respectively [60].

The development of both acceleration and sprinting mechanics can develop stride length and decreasing ground contact time, whilst optimizing ground reactive forces and decrease injury rates [61]. Faster athletes can produce a greater vertical impulse in a shorter ground contact time [62]. A further goal of maximal velocity sprinting is to achieve a high stride frequency combined with an optimal stride length [61, 62]. Furthermore, to mitigate against hamstring injuries during the terminal swing phase of sprinting, practitioners within elite female Gaelic team sports athletes are recommended to focus on multi-factorial approach which includes slow, high load eccentric contractions at the knee joint with development of efficient maximal velocity sprinting mechanics using front-side mechanics drilling constraints [63–66]. It is important that elite female Gaelic team sports athletes are exposed to adequate maximal velocity sprinting ($> 90\%$ of their maximal velocity) throughout the season which may provide a potential aid against hamstring injuries [65, 66]. Therefore, improving acceleration and maximal velocity mechanics could improve performance as well as reduce the risk of posterior-chain injuries across elite female Gaelic team sports athletes. Further research is needed to determine the effectiveness of acceleration and maximal velocity interventions in elite female Gaelic team sport athletes.

Power

Muscular power is an important attribute in determining athletic ability and predicting success in sport [4]. The countermovement jump (CMJ) is a widely used criterion and is an appropriate measure of muscular power in Gaelic team sport athletes [45]. Camogie players tend to have lower levels of jump performance (27.5 ± 3.1 cm) when compared to female Gaelic footballers despite having similar masses [43, 47]. In addition, the average reactive strength index (RSI) in elite intercounty Camogie

players was reported as 1.18 ± 0.21 [47]. In comparison to collegiate female Gaelic footballers in which the average RSI was reported as 1.22 ± 0.47 [67]. Additionally, in elite Irish female soccer players the average RSI was reported as 1.56 ± 0.25 [60]. Therefore, the implementation of reactive strength-based modalities in elite female Gaelic team sports athletes may lead to improvements in stretch shortening cycle ability. The CMJ performance of female Gaelic footballers was explored by Tucker and Reilly [45] with mean values of 44.9 ± 3.2 cm reported. These values are similar to data reported for female Australian football (45.5 ± 3.1 cm) players [44]. It has been demonstrated that long term (>10 weeks) plyometric training is an effective method to improve vertical performance jump in female athletes [68]. A 12-week plyometric training intervention combined with regular football training improved explosive strength and improvements transferred to soccer specific skills in female athletes [69]. Previous research in female team athletes supports the inclusion of plyometric training into a multi-faceted training program which can decrease knee valgus and increase ground reaction forces and increase hamstring strength [70–72]. It has been shown that combining plyometric or ballistic movement efforts with other resistance training methods (complex training) can lead to enhanced performance as opposed to plyometric training alone in female team athletes [57]. The utilization of a plyometric training stimulus could be beneficial to improve muscular power in elite female Gaelic team sport athletes. Therefore, future research is needed to determine the efficacy of different muscular power training modalities in elite female Gaelic team sports athletes. In conclusion, muscular power characteristics in elite female Gaelic team sport athletes are lower in comparison to their elite female team sports counterparts.

Injury incidence, risk and prevention

To embed successful injury management strategies in elite female Gaelic team sports, firstly one must understand the epidemiology of injury incidence and burden. Due to the high velocity, multi-directional, contact nature of Gaelic team sports, female athletes who participate in these sports, especially at an elite level, are inherently at risk of injury [73, 74]. In a self-reported participant injury survey across all levels of camogie, indicated that the most traumatic injury that occurred in the game/training activities was knee ligament damage (21% of all injuries) with over 60.8% of correspondences reporting lower limb injuries [3]. Furthermore, 85% of female correspondents felt obliged to continue playing whilst injured, reporting that they did not want to let their teammates down or believed they could manage the injury themselves [3]. Similarly, it was reported that lower limb injuries were

the most frequent in elite camogie (71.4%), with 23.8% of injuries occurring to thigh, knee (19%) and ankle (9.5%) [75]. This study reported that 19% of injuries were a result of sprinting, 14.3% resulting from a change of direction, 9.5% occurring from overloading and 4.8% occurring from a push off [75]. Although there is currently no data on injury rates in elite female Gaelic football, lower limb injuries were the most prevalent (67.09%) and caused the greatest injury burden (276.17 days absent per 1000 h) at a collegiate, sub-elite level [76]. Additionally, hamstring injuries were the most common (21.52%) followed by knee (12.66%), quadriceps (11.39%) and ankle (10.31%). It was reported that knee injuries cause the greatest injury burden (106.46 days absent per 1000 h) [76].

It has been reported that female team sports athletes are 3–6 times more likely to obtain an anterior cruciate ligament (ACL) injury than their male counterparts [77]. This high incidence has been largely attributed to anatomical and hormonal aspects including joint laxity, limb alignment, intercondylar notch proportions, ligament size and hormonal fluctuations [78]. Although there is no existing research on ACL injury rates in elite female Gaelic team sports athletes it is imperative that practitioners use different training modalities (resistance training, plyometrics, development of athletic motor skill competencies etc.) to support structural components and improve neuromuscular control of the knee joint and reduce the incidence of non-contact ACL injuries in field-based sports [54, 79]. It is reported during side cutting maneuvers in which internal tibia rotation occurs in <50 ms after initial contact in conjunction with reduced semitendinosus pre-activation, is associated with an increased risk of ACL injury [80]. Therefore, it may be worthwhile for elite Gaelic practitioners to consider specific training modalities to replicate forces produced <100 ms such as strength-speed, speed strength and velocity-based modalities [1, 2].

Developing stronger, resilient elite female Gaelic team sports athletes will enable them to master complex, athletic movements innate to the games and cope with the increased demands of training and competition without injury [2, 81]. An INT approach, which combines strength training, plyometrics and movement skill competencies to maximize female athletic performance [54, 55] has been recommended. This multifaceted training system can assist in the strengthening of tendons, ligaments and bones, enhanced balance and coordination between agonist and antagonist muscle groups and an improved capability to withstand training load from specific sports training and competition [53, 54]. It has been demonstrated that integrative neuromuscular training (INT) designed for female soccer players injury prevention simultaneously improved lower body performance

measures and improved movement mechanics [54, 55]. This accentuates the importance of a systematic IPP to decrease the severity of injuries [82]. An applied, minimally invasive example of an INT is the Gaelic specific injury prevention program (IPP) (GAA 15) [83–85]. Both LGFA and the Camogie Association have introduced the GAA specific GAA 15 IPP [84, 85]. The implementation of the GAA 15 IPP led to a 66% ($p=0.001$) reduction in training injuries, hamstring injuries, and non-contact injuries in a collegiate Gaelic team sports specific IPP intervention group [83]. A survey conducted on Camogie coaches and players reported that only 34% of coaches and 11.8% of players were using the GAA 15 and the Camogie Association should include a mandatory Camogie specific IPP concurrently with coach education programs [85]. The development of enhanced performance characteristics in conjunction with greater compliance of IPP's could mitigate against the injury risks associated with both elite female Gaelic team sports.

Anthropometric characteristics

The anthropometric characteristics of athletes are an important health and performance variable [86]. The information is beneficial for coaches and practitioners to inform training practices and nutritional strategies as well as potentially guide playing positions. Currently, there is a paucity of anthropometric data on elite female Gaelic footballers compared to camogie players, as demonstrated in Table 1. Players ranged between 18–29 years of age, 1.58–1.74 m in stature, 56.4–75.9 kg in body mass and had a body fat percentage of 21–26%. The age, stature and mass range of female Gaelic footballers are comparable to other female team sports such as rugby, handball, and hockey [86–88], however, the range observed in female soccer players is far greater and this highlights the heterogeneity among top level soccer players [14]. When comparing the data from earlier studies [43] to the most recent [24, 25, 29, 89] players' stature and body mass have remained similar despite the sport evolving in recent years and the advent of strength and

conditioning programs. Positional differences of female Gaelic footballers have been assessed in one study however, this was carried out on sub-elite players [90]. The authors reported similarities between positions for both body mass and body mass index (BMI), but a ~3–7 cm mean difference in stature was observed between midfielders and other positions. However, it is unclear if these positional differences exist at an elite level.

To date, only one study [43] has reported body fat percentage values for elite female Gaelic footballers using a 4-site skinfold assessment method. The body fat percentage values reported by Keane et al. [43] were less than the non-exercising controls in the study but are also comparable to elite female handball [87] and hockey players [88]. In contrast, female soccer players tend to have a lower body fat percentage with ranges of 15–22% reported [14] whilst elite rugby league players have slightly higher values typically ranging from 23–39% depending on playing position [91]. Body composition data pertaining to female GAA players requires updating before positional differences can be explored. Although not Gaelic team sports specific, there are best practice recommendations in terms of assessing body composition although this will be dependent on what equipment is available. Where feasible, DEXA or skinfolds (ISAK accredited) should be used [92].

Nutritional considerations

Nutrition plays a significant role in athletic performance, achieved through the dietary influence on body composition, training adaptation, and recovery [93, 94]. There has been growing interest in the nutritional requirements of Gaelic team sport athletes and subsequently, a number of recent nutritional studies have been conducted. However, to date many of these studies and subsequent recommendations are based on male athletes and as such there is an absence of sex specific nutrition guidelines. This is a vital first step in promoting the health and performance of elite female Gaelic team sport athletes.

Table 1 The anthropometric characteristics of elite female Gaelic team sport players

Reference	Country	Sport	Standard	Age (years)	Stature (m)	Body mass (kg)	% Body fat
O'Grady et al. [29] $n=43$	Ireland	Camogie	Elite	23 ± 5.0	1.74 ± 0.05	68 ± 9	–
Connors et al. [47] $n=45$	Ireland	Camogie	Elite	23.3 ± 3.47	1.68 ± 0.06	68.4 ± 7.4	–
Renard et al. [89] $n=328$	Ireland	Camogie, GF		23.7 ± 3.0	1.66 ± 0.08	65.8 ± 9.1	–
Young et al. [24] $n=36$	Ireland	Camogie	Elite	23 ± 4.0	1.68 ± 0.05	65 ± 6.0	–
Connors et al. [25] $n=24$	Ireland	Camogie	Elite	23.5 ± 3.37	1.68 ± 0.06	68.5 ± 7.4	–
Buckley and Blake [75] $n=62$	Ireland	Camogie	Elite	22.9 ± 3.6	1.68 ± 0.06	64.8 ± 6.3	–
Keane et al. [43] $n=46$	Ireland	GF	Elite	18–29	1.66 ± 0.05	63.7 ± 7.3	23.3 ± 2.3

Energy and macronutrient intake

Research has consistently shown sub-optimal dietary practices amongst elite and sub-elite male Gaelic team sport athletes, most notably failure to meet both energy and carbohydrate requirements. Gaelic football players dietary intake has been identified as inadequate to meet recommendations, with average energy deficits of 12.3% per day [95], carbohydrate intakes of 3.4–3.7 g kg day⁻¹ [95–97] lower than minimum recommendations of 5 g kg day⁻¹. It remains to be established if similar dietary intakes are adopted by female Gaelic team sport athletes, however similar findings have been reported with other female field-based team sport athletes [98–100]. Not only does an inadequacy of energy and carbohydrate intake in a team sport context limit substrate availability during training and matches as well as impair recovery [101], if athletes have inadequate energy intake to match exercise energy expenditure, they are described as having low energy availability (LEA) [102] or relative energy deficiency in sport (RED-S) [103]. Growing evidence indicates that unhealthy dietary behaviors and negative body image issues are common in both male and female athletes, but it is still widely accepted that female athletes are more likely to experience LEA. The underlying cause of RED-S is LEA and is calculated by subtracting exercise energy expenditure (EEE), kcal/d from tkcal/d. If the RED-S syndrome is present in an athlete, either inadvertently or through purposeful dieting or disordered eating, athletes can experience increased fatigue, injuries or illness, nutrient deficiencies, menstrual dysfunction, poor bone health, and lack of improvement in performance. In addition, athletes can experience impairments in metabolic rate, immunity, protein synthesis and cardiovascular health [103]. To date, there is limited research into the prevalence of LEA in female Gaelic team sport athletes' players. One study examining the prevalence of LEA in athletic and recreationally active females in Ireland, including 118 Gaelic footballers and 45 camogie players, found 40% (39.7%, n=331) of participants were classified as being 'at risk' of LEA according to the Low Energy Availability for Females Questionnaire scoring system [104]. Furthermore, risk was 1.7 and 1.8 times more likely to occur among elite (intercounty) athletes compared to those who were recreationally active. Further research on energy availability in elite female Gaelic team sports is needed to consolidate these findings.

On the contrary, elite and sub-elite male Gaelic team sport athletes have been shown to exceed protein and fat recommendations with protein intakes of 1.9–2.1 g kg day⁻¹ [96, 97, 105] towards the higher end of recommendations and fat intakes of 31–37.5% of total daily energy intake (TDEI) also at the higher end of recommendations [106]. Over consumption of protein and

fat has previously been associated with reduced carbohydrate intake in elite male soccer players [107] and may partly account for the inadequate carbohydrate intake observed in male Gaelic team sport athletes. On the contrary, it has been suggested the female athletes are more likely to be at risk for low protein intakes [108]. Currently it is not known if this high fat and high or low protein eating behaviors are seen in female Gaelic team sport athletes and this warrants further investigation.

Micronutrient intake

With regards to micronutrient intake, active females often report low levels of micronutrients especially in cases where energy intake is being restricted, food groups are eliminated, a special diet is followed (e.g., vegan, vegetarian, or fad diet), the athlete has an eating disorder, or there are health issues that alter nutrient absorption [108–110]. Inadequate intakes of calcium and iron are common in females, with inadequate iron intakes previously documented in female team sports [111]. In Ireland, vitamin D insufficiency and deficiency is pervasive in athletes, and previous studies have identified a particularly high prevalence of vitamin D insufficiency/deficiency in Gaelic footballers [112, 113]. To date, there has only been one study (abstract only) that has investigated the dietary intakes of calcium, iron, and vitamin D in sub-elite (n=23) and elite (n=26) female Gaelic footballers [114]. Overall, median intakes of calcium, iron and vitamin D in both sub-elite and elite footballers were lower than the respective recommended daily intake (RDA) [114]. In addition, reported iron intakes of sub-elite players were significantly lower than elite players ($p < 0.05$), with no differences in calcium and vitamin D intakes between sub-groups. Interestingly, a minority of elite players (n=2) in this study met the RDAs for calcium, iron, and vitamin D with the aid of dietary supplements [114]. A recent addition to the literature investigating the sex differences in prevalence of dietary supplement use in elite athletes, participating in individual and team sports reported that supplement use was slightly higher in males than in females however female athletes appeared to consume more iron supplements and supplements intended to improve health [115]. To date, there has been no research into supplement usage amongst female or male Gaelic team sport athletes.

Hydration

The detrimental effects of dehydration on performance have been well documented including, an increase in core body temperature, cardiovascular strain and glycogen utilization and impaired cognitive performance [116, 117]. Despite the obvious benefit of starting exercise in a euhydrated state, many athletes begin exercise

already dehydrated [118]. There is a dearth of literature on the hydration status of athletes in Ireland; the majority having focused on men's Gaelic football [119, 120]. However, one study that included sub-elite female Gaelic team sport athletes found that most participants were euhydrated before and after exercise [112]. This study reported that 25% of the cohort were dehydrated pre and post exercise which is lower than their male counterparts (35 and 48% respectively). This may be in part explained by the lower sweat rates in female athletes [116] or the reliance on percentage body weight loss as criterion for dehydration post-exercise [120]. In contrast, an earlier study by Newell and colleagues reported only 10% of elite Gaelic football players (all males) commenced exercise in a dehydrated state [120]. Further research with elite male and female Gaelic players using validated measures both pre and post exercise are needed.

Nutritional knowledge

Nutritional knowledge has been shown to play a significant role in adopting optimal nutrition practices, such as sufficient macro- and micronutrient intake. A previous investigation reported inadequate levels of nutrition knowledge in female Gaelic team sport players (football and camogie). It was reported that their mean nutrition knowledge score was $46.0\% \pm 11.8\%$ and classified as 'poor' [89]. Players scored better on food-based questions yet lacked knowledge surrounding specific macronutrient recommendations. Interestingly, elite players scored greater ($+4.5\text{--}5.9\%$, $p < 0.05$) than sub elite players. Players with higher levels of general education, history of formal nutrition education and previous advice from a nutritionist also presented greater nutrition knowledge ($+3.7\text{--}7.5\%$, $p < 0.05$). The poor nutrition knowledge of female Gaelic team sports players may have negative consequences for their dietary behavior. Despite a current lack of dietary investigation within the population, other female team sport athletes with poor nutritional knowledge have been shown have insufficient energy and carbohydrate intake [98, 99]. Similarly, this relationship between poor nutritional knowledge and sub-optimal dietary intake has been seen in male Gaelic football players [95, 97]. As such it would be beneficial to measure dietary intake and assess nutritional knowledge simultaneously in elite female Gaelic team sport athletes to see if knowledge translates to practice. In addition, focus should be placed on strategies to enhance nutrition knowledge of sport-specific concepts, addressing macro and micronutrient recommendations as players scored poorest on these aspects.

Female physiology

The menstrual cycle (MC)

The MC results in large variations in both oestrogen and progesterone concentrations [121]. Although there is no research available on elite female Gaelic team athletes, these variations in reproductive hormones may influence injury risk and performance. The effect of estrogen has been suggested as a possible injury mechanism for ACLs in female athletes [122], however there is a lack of agreement regarding the influence of sex hormones on ACL injury rates. It has been reported that female soccer players had an increased incident rate during the luteal phase of the MC [123] Martin et al. [124] reported that injury rates were 47% and 32% greater in the late follicular phase in comparison to the early follicular and luteal phase. However, caution is needed in the interpretation of these studies as the methodological process (small sample size), ecological validity and terminology (phase definition and confirmation) may have affected the interpretation of the original results.

As females train and compete during all stages of their MC, the potential impact on performance is of worthy consideration. There has been some research conducted in the area with conflicting findings to date and no consensus on whether performance is affected by MC phases [125, 126]. A recent meta-analysis showed that exercise performance may be trivially reduced during the early follicular phase of the MC, compared to other phases [125]. However, it must be noted that there was large between-study variation and as with most research in this area, participants were studied at a group level. As it is commonly known the effects of the menstrual cycle are highly individualized, it is recommended that a personalized approach should be taken whereby players track their own cycles and any perceived phase-related effects on performance should be noted.

Hormonal contraceptives

There is a plethora of hormonal contraceptives available including oral contraceptives (OC), implants, injections patches, and intra-uterine systems [127]. OCs appear to be the most popular with Martin et al. reporting that 49.5% of 430 elite female athletes surveyed from 24 different sports were currently using OC and 69.8% had used the OC previously. This study also revealed that female athletes reported the ability to regulate menstruation during competition/training as a positive effect and reason for using OCs [127]. To date, there has been no research conducted on the use of hormonal contraceptives in elite female Gaelic team sports players or the reasons for initiation. Furthermore, there are conflicting reports on the effects of OC use on exercise performance throughout the literature [128, 129], with the most recent

meta-analysis reporting that OC use might result in slightly inferior exercise performance when compared to naturally menstruating women, however any affect is most likely trivial [129]. Further research is necessary to determine whether the use of OC has positive, negligible, or detrimental effects on female athletic performance with regards to all sports [14].

Limitations and future recommendations

It is important to note the limitations of the current review and in the literature. A narrative style of review was chosen as it was not possible to adopt a systematic approach due to the paucity of research currently available in the field of elite female Gaelic team sport players. In addition, this literature review focused on seven key research topics that are of pertinent interest to researchers and practitioners working in Games, nonetheless, there are other performance parameters that have not been discussed, due to the lack of existing literature. Finally, some of the studies included in the review are of mediocre quality and often include small sample sizes, as such results need to be interpreted with some degree of caution. This review has summarized the available research, as well as highlighting knowledge gaps and areas of future research needs within the domain of elite female Gaelic teams sports. These areas for future research have been highlighted in the individual sections, with a summary provided in Fig. 1.

Summary and conclusion

Interest in female athletes has grown exponentially over the past decade, with recent research efforts focusing on the match demands, physiological characteristics as well as specific health and performance considerations in female intermittent team-based sports. To the best of the authors knowledge, this is the first review summarizing the current research available in elite female Gaelic team sports, with a focus on seven specific areas including: physical and physiological demands, anthropometric and performance characteristics, injury risk, nutritional considerations, and female physiology.

In summary, there appears to be unique physical demands data in match play across positions in camogie with midfielders having greater total and sprint distance decrements. There is currently no comparative data available in ladies Gaelic football. Similarly, there is no research available on the physiological demands of both elite female Gaelic team sports. According to existing literature, performance characteristics such as speed and power are lower in this population compared to other elite female team sports. Based on the research available, the anthropometric characteristics of elite female Gaelic team sport players appear homogenous with some positional differences in stature observed at a sub-elite level. Previous research has demonstrated a high prevalence of lower limb injuries in female elite Gaelic team sports with 85% of players reporting they




Fig. 1 Future research directions in elite female Gaelic team sports

felt obliged to continue playing whilst injured. The provision of quality, evidence-based strength & conditioning such as the Gaelic specific IPP could help mitigate these injury risks with positive results already observed at a sub-elite level. With regards to nutrition, female Gaelic team sport players have been shown to have inadequate nutrition knowledge which may put them at a higher risk of LEA. In addition, previous research has shown sub-optimal intakes of certain micronutrients including vitamin D, iron, and calcium in this population. Positively, it would appear female Gaelic team sport players adopt appropriate hydration practices. Finally, although menstrual cycle phase and oral contraceptives have been shown to influence performance in other female intermittent sports, to date there has not been any research carried out in elite female Gaelic team sport players.

To conclude, the current work has summarized the existing literature in the area and created a roadmap for researchers in highlighting the existing knowledge gaps that need to be addressed in this field. Addressing these, would help inform future recommendations and make it easier for practitioners to adopt an evidence-based approach when working with elite female Gaelic team sport players.

Abbreviations

AFL: Australian Rules Football; GPS: Global position system; m: Meters; m min^{-1} : Meters per minute; km h^{-1} : Kilometers per hour; HR: Heart rate; HR_{max} : Maximum heart rate; HR_{peak} : Heart Rate peak; b min^{-1} : Beats per minute; $\text{mL kg}^{-1} \text{min}^{-1}$: Milliliters per kilogram per minute; IMTP: Isometric mid-thigh pull; N: Newtons; N/kg: Relative peak force; RFD: Rate of force development; s: Seconds; CMJ: Counter-movement jump; RSI: Reactive strength index; ACL: Anterior cruciate ligament; ms: Milliseconds; INT: Integrated neuromuscular training; IPP: Injury prevention program; GAA: Gaelic Athletic Association; LGFA: Ladies Gaelic Football Association; BMI: Body mass index; DEXA: Dual-energy X-ray absorptiometry; ISAK: International Society for the Advancement of Kinanthropometry; LEA: Low energy availability; REDS: Relative energy deficiency in sport; EEE: Exercise energy expenditure; Kcal/d: Kilocalorie; TDEI: Total daily energy intake; RDA: Recommended daily allowance; MC: Menstrual cycle; OC: Oral contraceptives; RPE: Rate of perceived exertion.

Acknowledgements

Not applicable.

Author contributions

JD, KC & KK: conceived the manuscript idea. JD: performed the initial search and data extraction, drafted the article and made critical revisions, read and approved the final manuscript. KC: drafted the article and made revisions, read and approved the final manuscript. KK: drafted the article and made critical revisions, read and approved the final manuscript. All authors read and approved the final manuscript.

Funding

No funding was obtained for this project.

Availability of data and materials

Not applicable.

Declarations

Ethics approval and consent to participate

Not applicable.

Consent for publication

Not applicable.

Competing interests

The authors declare that they have no competing interests.

Author details

¹Department of Sport, Exercise & Nutrition, School of Science & Computing, Atlantic Technological University Galway, Dublin Road, Galway, Ireland. ²Gaelic Sports Research Centre, Technological University Dublin-Tallaght Campus, Tallaght, Dublin 24, Dublin, Ireland.

Received: 27 October 2021 Accepted: 22 August 2022

Published online: 02 September 2022

References

- Duggan JD, Moody JA, Byrne P, Ryan L. Strength & conditioning recommendations for female GAA athletes: the Camogie player. *Strength Cond J*. 2020;42(4):105–24.
- Duggan JD, Moody JA, Byrne P, McGahan J, Kirszenstein L. Considerations and guidelines on athletic development for youth Gaelic Athletic players. *Strength Cond J*. 2021. <https://doi.org/10.1519/SSC.0000000000000638>.
- O'Connor S, Leahy R, Whyte E, O'Donovan P, Fortington L. Understanding injuries in the Gaelic sport of camogie: the first national survey of self-reported worst injuries. *Inter J Ath Ther Train*. 2019;24(6):243–8.
- Reilly T, Collins K. Science and the Gaelic sports: Gaelic Football and Hurling. *Eur J Sport Sci*. 2008;8(5):231–40.
- Beasley KJ. Nutrition and Gaelic football: Review, recommendations, and future considerations. *Int J Sport Nutr Exerc Metab*. 2015;25(1):1–13.
- Kelly G, McKenna O, Courtney S, Collins K, Bradley J, Martin D. Benchmarking successful performances in elite Ladies Gaelic football. *Int J Perform Anal Sport*. 2022;22(7):51–65. <https://doi.org/10.1080/24748668.2021.2020031>.
- Heyward O, Nicholson B, Emmonds S, Roe G, Jones B. Physical preparation in female rugby codes: an investigation of current practices. *Front Sports Act Living*. 2020. <https://doi.org/10.3389/fspor.2020.584194>.
- Nimphius S. Exercise and sport science failing by design in understanding female athletes. *Int J Sports Physiol Perform*. 2019. <https://doi.org/10.1123/ijspp.2019-0703>.
- Cowley E, Olenick AA, McNulty K, Ross, EZ. "Invisible Sportswomen" The sex data gap in sport and exercise science research. *Women Sports Phys Act*. 2021. <https://doi.org/10.1123/wspaj.2021-0028>.
- Costello JT, Bieuzen F, Bleakley CM. Where are all the female participants in sports and exercise medicine research? *Eur J Sport Sci*. 2014;14(8):847–51.
- Heyward O, Emmonds S, Roe G, Scantlebury S, Stokes K, Jones B. Applied sport science and medicine of women's rugby codes: a systematic-scoping review and consensus on future research priorities protocol. *BMJ Open Sport Exerc Med*. 2021;7(3):e001108. <https://doi.org/10.1136/bmjsem-2021-001108>.
- Emmonds S, Heyward O, Jones B. The challenge of applying and undertaking research in female sport. *Sports Med Open*. 2019;5(1):1–4.
- Datson N, Hulton A, Andersson H, Lewis T, Weston M, Drust B, Gregson W. Applied physiology of female soccer: an update. *Sports Med*. 2009;44(9):1225–40.
- Randall RK, Clifford T, Drust B, Moss SL, Unnithan VB, De Ste Croix MBA, Datson N, Martin D, Mayho H, Carter JM, Rollo I. Physiological characteristics of female soccer players and health and performance considerations: a narrative review. *Sports Med*. 2021;51(7):1377–99.
- Sella FS, McMaster DT, Beaven CM, Gill ND, Hébert-Losier K. match demands, anthropometric characteristics, and physical qualities of

- female rugby sevens athletes: a systematic review. *J Strength Cond Res.* 2019;33(12):3463–74.
16. Clarke A, Ryan S, Couvalias G, Dascombe BJ, Coutts AJ, Kempton T. Physical demands and technical performance in Australian football league women's (AFLW) competition match-play. *J Sci Med Sport.* 2018;21(7):748–52.
 17. Cummins C, Orr R, O'Connor H. Global positioning systems (GPS) and microtechnology sensors in team sports: a systematic review. *Sports Med.* 2013;43(10):1025–42.
 18. Dellaserra CL, Gao Y, Ransdell L. Use of integrated technology in team sports: a review of opportunities, challenges, and future directions for athletes. *J Strength Cond Res.* 2014;28(2):556–73.
 19. Malone S, Collins K. Effect of game design, goal type, and player numbers on the physiological and physical demands of hurling-specific small-sided games. *J Strength Cond Res.* 2017;31(6):1493–9.
 20. Malone S, Solan B, Collins K. The running performance profile of elite Gaelic football match-play. *J Strength Cond Res.* 2017;31(1):30–6.
 21. McGahan JH, Mangan S, Collins K, Burns C, Gabbett TJ, O'Neill C. Match-play running demands and technical performance among elite Gaelic footballers: Does divisional status count? *J Strength Cond Res.* 2021;35(1):169–75.
 22. Young D, Mouro L, Beato M, Coratella G. The match heart rate and running profile of elite under-21 Hurlers during competitive match-play. *J Strength Cond Res.* 2018;32(10):2925–33.
 23. Young D, Beato M, Mouro L, Coratella G. Match-play temporal and position-specific physical and physiological demands of senior hurlers. *J Strength Cond Res.* 2020;34(6):1759–68.
 24. Young D, O'Grady M, Coratella G. The match-play running performance of elite Camogie players across halves of play. *Sports Sci Health.* 2020. <https://doi.org/10.1007/s11332-020-00672-0>.
 25. Connors P, Earls D, Browne D, Fitzpatrick P, Rankin P. The positional demands of inter-county Camogie. *Sport Sci Perform Rep.* 2020. https://sportperfsci.com/wp-content/uploads/2020/07/SPSR110_Connors.pdf.
 26. Connors P, Earls D, Browne D, Fitzpatrick P, Rankin P. The positional and temporal demands of elite inter-county camogie match play across 5-intervals. *Sports Sci Health.* 2022. <https://doi.org/10.1007/s11332-022-00904-5>.
 27. Novak AR, Impellizzeri FM, Trivedi A, Coutts AJ, McCall A. Analysis of the worst-case scenarios in an elite football team: towards a better understanding and application. *J Sports Sci.* 2021;39(16):1850–9.
 28. McGuinness A, Passmore D, Malone S, Collins K. Peak running intensity of elite female field hockey players during competitive match play. *J Strength Cond Res.* 2020. <https://doi.org/10.1519/JSC.00000000000003582>.
 29. O'Grady M, Young D, Collins K, Keane J, Malone S, Coratella G. An investigation of the sprint performance of senior elite camogie players during competitive play. *Sports Sci Health.* 2022. <https://doi.org/10.1007/s11332-021-00874-0>.
 30. Callanan D, Rankin P, Fitzpatrick P. An analysis of the game movement demands of women's interprovincial rugby union. *J Strength Cond Res.* 2021. <https://doi.org/10.1519/JSC.0000000000004065>.
 31. McErlean C, Cassidy J, O'Donoghue P. Time-motion analysis of gender and positional effects on work-rate in elite Gaelic football competition. *J Hum Mov Studies.* 2000;2000(38):269–86.
 32. Duggan JD, Moody JA, Byrne PJ, Cooper S-M, Ryan L. Training load monitoring considerations for female Gaelic team sports: from theory to practice. *Sports.* 2021;9(6):84. <https://doi.org/10.3390/sports9060084>.
 33. McFadden BA, Walker AJ, Bozzini BN, Sanders DJ, Arent SM. Comparison of internal and external training loads in male and female collegiate soccer players during practices vs. games. *J Strength Cond Res.* 2020;34(4):969–74.
 34. Busbridge AR, Hamlin MJ, Jowsey JA, Vanner MH, Olsen PD. Running demands of provincial women's rugby union matches in New Zealand. *J Strength Cond Res.* 2020. <https://doi.org/10.1519/JSC.00000000000003579>.
 35. Andersson H, Andersson HÅ, Randers MB, Heiner-Møller A, Krstrup P, Mohr M. Elite female soccer players perform more high-intensity running when playing in international games compared with domestic games. *J Strength Cond Res.* 2010;24(4):912–9.
 36. Bradley PS, Vescovi JD. Velocity thresholds for women's soccer matches: Sex specificity dictates high-speed running and sprinting thresholds-Female athlete in motion (FAIM). *Int J Sports Physiol Perform.* 2015;10(1):112–6.
 37. Trewin J, Meylan C, Varley MC, Cronin J. The match-to-match variation of match-running in elite female soccer. *J Sci Med Sport.* 2018;21(2):196–201.
 38. Krstrup P, Mohr M, Ellingsgaard H, Bangsbo J. Physical demands during an elite female soccer game: importance of training status. *Med Sci Sports Exerc.* 2005;37(7):1242–8.
 39. McGuinness A, Malone S, Petrakos G, Collins K. Physical and physiological demands of elite international female field hockey players during competitive match play. *J Strength Cond Res.* 2019;33:3105–13.
 40. Mara JK, Thompson KG, Pumpa KL. Physical and physiological characteristics of various-sided games in elite women's soccer. *Int J Sports Physiol Perform.* 2016;11(7):953–8.
 41. Mangan S, Collins K, Burns C, O'Neill C. A tactical periodization model for Gaelic football. *Int J Sports Sci Coach.* 2021. <https://doi.org/10.1177/17479541211016269>.
 42. Dekkers T, O'Sullivan K, Blake C, McVeigh JG, Collins K. Epidemiology, and moderators of injury in Gaelic football: a systematic review and meta-analysis. *J Sci Med Sport.* 2021;14:51440–2440(21)00455–2.
 43. Keane A, Scott MA, Dugdill L, Reilly T. Fitness test profiles as determined by the Eurofit test battery in elite female Gaelic football players. *J Strength Cond Res.* 2010;24(6):1502–6.
 44. Wyckelsma V, Aughey R, McKenna M. Physiological responses and movement demands of elite women playing Australian football in the midfield position. *J Sci Med Sport.* 2010;13(1):15–6.
 45. Tucker L, Reilly T. Physiological and anthropometric characteristics of female Gaelic football players. *Science and Football V: Proceedings of the Fifth World Congress of Science and Football, 2005:* 27–30.
 46. McIntyre MC. A comparison of the physiological profiles of elite Gaelic footballers, hurlers, and soccer players. *Br J Sports Med.* 2004;39(7):437–9.
 47. Connors PM, Browne DT, Earls D, Fitzpatrick P, Rankin P. The physical characteristics of elite camogie players. *J Sports Med Phys Fitness.* 2021. <https://doi.org/10.23736/S0022-4707.21.12429-6>.
 48. Yao X, Curtis C, Turner A, Bishop C, Austerberry A, Chavda S. Anthropometric profiles and physical characteristics in competitive female english premiership rugby union players. *Int J Sports Physiol Perform.* 2021. <https://doi.org/10.1123/ijsp.2020-0017>. Epub ahead of print.
 49. Suchomel TJ, Nimphius S, Bellon CR, Stone MH. The importance of muscular strength: training considerations. *Sports Med.* 2018;48(4):765–85.
 50. Suchomel TJ, Nimphius S, Stone MH. The importance of muscular strength in athletic performance. *Sports Med.* 2016;46(10):1419–49.
 51. Myer GD, Ford KR, Hewett TE. Methodological approaches and rationale for training to prevent anterior cruciate ligament injuries in female athletes. *Scand J Med Sci Sports.* 2004;14(5):275–85.
 52. Myer GD, Ford KR, Hewett TE. Rationale and clinical techniques for anterior cruciate ligament injury prevention among female athletes. *J Ath Train.* 2004;39(4):352–64.
 53. Myer GD, Paterno MV, Ford KR, Hewett TE. Neuromuscular training techniques to target deficits before return to sport after anterior cruciate ligament reconstruction. *J Strength Cond Res.* 2004;22(3):987–1014.
 54. Myer GD, Ford KR, Palumbo JP, Hewett TE. Neuromuscular training improves performance and lower-extremity biomechanics in female athletes. *J Strength Cond Res.* 2005;19(1):51–60.
 55. Myer GD, Brent JL, Ford KR, Hewett TE. Real time assessment and neuromuscular training feedback techniques to prevent anterior cruciate ligament injury in female athletes. *Strength Cond J.* 2011;33:21–35.
 56. Chilibeck PD, Calder AW, Sale DG, Webber CE. A comparison of strength & muscle mass increases during resistance training in young women. *Eur J Appl Physiol.* 1998;77:170–5.
 57. Grieco CR, Cortes N, Greska EK, Lucci S, Onate JA. Effects of a combined resistance-plyometric training program on muscular strength, running economy, and VO₂peak in division I female soccer players. *J Strength Cond Res.* 2012;26(9):2570–6.
 58. Strudwick A, Doran D, Reilly T. Anthropometric and fitness profiles of elite players in two football codes. *J Sports Med Fit.* 2002;42(2):233–42.
 59. Reilly T, Doran D. Science and Gaelic football: a review. *J Sports Sci.* 2001;19(3):181–93.

60. Doyle B, Browne D, Horan D. Differences in anthropometric and physical performance characteristics between U17, U19 and senior Irish female international football players. *Int J Sports Sci Coach*. 2021;16(2):352–9.
61. Weyland PG, Davies JA. Running performance has a structural basis. *J Exp Biol*. 2005;208(14):2625.
62. Weyland PG, Sternlight DB, Belizzi MJ, Wright S. Faster top running speeds are achieved with greater ground forces not more rapid leg movements. *J Appl Physiol*. 2000;89(5):1991–9.
63. Guex K, Millet GP. Conceptual framework for strengthening exercises to prevent hamstring strains. *Sports Med*. 2013;43(12):1207–15.
64. Mendiguchia J, Martínez-Ruiz E, Edouard P, Morin JB, Martínez-Martinez F, Idoate F, Mendez-Villanueva A. A multifactorial, criteria-based progressive algorithm for hamstring injury treatment. *Med Sci Sports Exerc*. 2017;49(7):1482–92.
65. Mendiguchia J, Castano-Zambudio A, Jimenez-Rayes P, Morin JB, Edouard P, Conceicao F, Dodo J, Colyer S. Can we modify maximal speed running posture? Implications for performance and hamstring injuries management. *Int J Sport Physiol Perform*. 2021. <https://doi.org/10.1123/ijspp.2021-0107>.
66. Lahti J, Mendiguchia J, Ahtiainen J, et al. Multifactorial individualised programme for hamstring muscle injury risk reduction in professional football: protocol for a prospective cohort study. *BMJ Open Sport & Exercise Medicine*. 2020;6:e000758. <https://doi.org/10.1136/bmjsem-2020-000758>.
67. Comyns TM, Flanagan EP, Fleming S, Fitzgerald E, Harper DJ. Inter-day reliability and usefulness of a reactive strength index derived from 2 maximal rebound jump tests. *Int J Sports Physiol Perform*. 2019;29:1200–4.
68. Stojanović E, Ristić V, McMaster DT, Milanovic Z. Effect of plyometric training on vertical jump performance in female athletes: a systematic review and meta-analysis. *Sports Med*. 2017;47(5):975–86.
69. Campo SS, Vaeynes R, Philippaerts RM, Redondo JC, De Benito AM, Cuadrado G. Effects of lower-limb plyometric training on body composition, explosive strength, and kicking speed in female soccer players. *J Strength Cond Res*. 2009;23(6):1714–22.
70. Hewitt TE, Stroupe AL, Nance TA, Noyes FR. Plyometric training for female athletes decreased impact forces and increased hamstring torques. *Am J Sports Med*. 1996;24(6):765–73.
71. Chimera NJ, Swanik KA, Swanik CB, Straub SJ. Effects of plyometric training on muscle-activation strategies and performance in female athletes. *J Athl Train*. 2004;39(1):24–31.
72. Hewitt TE, Myer GD, Ford KR. Reducing knee and anterior cruciate ligament injuries amongst female athletes. *J Knee Surg*. 2005;18(1):82–8.
73. Murphy JC, O'Malley E, Gissane C, Blake C. Incidence of injury in Gaelic football: a 4-year prospective study. *Am J Sports Med*. 2012;40(9):2113–20.
74. Roe M, Blake C, Gissane C, Collins K. Injury scheme claims in Gaelic games: a review of 2007–2014. *J Athl Train*. 2016;51(4):303–8.
75. Buckley CS, Blake C. The incidence of injury in elite camogie, an in-season prospective study. *BMJ Open Sport Exerc Med*. 2018;4(1):e000315. <https://doi.org/10.1136/bmjsem-2017-000315>.
76. O'Connor S, Bruce C, Teahan C, McDermott E, Whyte E. Injuries in collegiate ladies Gaelic footballers: a 2-season prospective cohort study. *J Sport Rehab*. 2020;30(2):261–6.
77. Voskanian N. ACL Injury prevention in female athletes: review of the literature and practical considerations in implementing an ACL prevention program. *Curr Rev Musculoskelet Med*. 2013;6(2):158–63.
78. Hewitt TE. Neuromuscular and hormonal factors associated with knee injuries in female athletes: strategies for intervention. *Sports Med*. 2000;29(5):313–27.
79. Alentorn-Geli E, Myer GD, Silvers HJ, Samitier G, Romero D, Lázaro-Haro C, Cugat R. Prevention of non-contact anterior cruciate ligament injuries in soccer players. Part 2: a review of prevention programs aimed to modify risk factors and to reduce injury rates. *Knee Surg Sports Traumatol Arthrosc*. 2009;17(8):859–79.
80. Zebis MK, Aagaard P, Andersen LL, Hölmich P, Clausen MB, Brandt M, Husted RS, Lauridsen HB, Curtis DJ, Bencke J. First-time anterior cruciate ligament injury in adolescent female elite athletes: a prospective cohort study to identify modifiable risk factors. *Knee Surg Sports Traumatol Arthrosc*. 2021. <https://doi.org/10.1007/s00167-021-06595-8>.
81. Taberner M, van Dyk N, Allen T, et al. Physical preparation and return to performance of an elite female football player following ACL reconstruction: a journey to the FIFA Women's World Cup. *BMJ Open Sport & Exerc Med*. 2020;6(1):e000843. <https://doi.org/10.1136/bmjsem-2020-000843>.
82. Laursen JB, Bertelsen DM, Andersen LB. The effectiveness of exercise interventions to prevent sports injuries: a systematic review and meta-analysis of randomised controlled trials. *Br J Sports Med*. 2014;48(11):871–7.
83. Schlingermann BE, Lodge CA, Gissane C, Rankin PM. Effects of the Gaelic Athletic Association 15 on lower extremity injury incidence and neuromuscular functional outcomes in collegiate Gaelic games. *J Strength Cond Res*. 2018;32(7):1993–2001.
84. O'Connor S, Lacey P. Can we improve coaches' injury prevention views and implementation practices in the community female Gaelic sport of camogie? *BMJ Open Sport Exerc Med*. 2020;6(1):e000732. <https://doi.org/10.1136/bmjsem-2019-000732>.
85. O'Connor S, Whyte E, O'Hanlon S, Fortington L. Coaches and players views towards injury prevention exercise programs in Camogie: A cross sectional survey. *Athl Train Sports Health Care*. 2021;13(4):e202-211.
86. Ackland TR, Lohman TG, Sundgot-Borgen J, Maughan RJ, Meyer NL, Stewart AD, et al. Current status of body composition assessment in sport: review and position statement on behalf of the ad hoc research working group on body composition health and performance, under the auspices of the IOC. *Medical Commission*. *Sports Med*. 2012;42(3):227–49.
87. Milanese C, Piscitelli F, Lampis C, Zancanaro C. Anthropometry and body composition of female handball players according to competitive level or the playing position. *J Sports Sci*. 2011;29(12):1301–9. <https://doi.org/10.1080/02640414.2011.591419>.
88. Keogh JW, Weber CL, Dalton CT. Evaluation of anthropometric, physiological, and skill-related tests for talent identification in female field hockey. *Can J Appl Physiol*. 2003;28(3):397–409. <https://doi.org/10.1139/h03-029>.
89. Renard M, Kelly DT, Ni Chéilleachair N, O Cathain C. Evaluation of nutrition knowledge in female Gaelic games players. *Sports*. 2020;8(12):154. <https://doi.org/10.3390/sports8120154>.
90. Brown J, Papadopoulos C, Pritchett R. Examination of Injury in Female Gaelic Football. *Int J Exerc Sci*. 2013;6(2):98–105.
91. Jones B, Emmonds S, Hind K, Nicholson G, Rutherford Z, Till K. Physical qualities of International female rugby league players by playing position. *J Strength Cond*. 2015;30(5):1333–40.
92. Kasper AM, Langan-Evans C, Hudson JF, Brownlee TE, Harper LD, Naughton RJ, Morton JP, Close GL. Come back skinfolds, all is forgiven: a narrative review of the efficacy of common body composition methods in applied sports practice. *Nutrients*. 2021;13(4):1075. <https://doi.org/10.3390/nu13041075>.
93. Garthe I, Raastad T, Refsnes PE, Sundgot-Borgen J. Effect of nutritional intervention on body composition and performance in elite athletes. *Eur J Sports Sci*. 2013;13(3):295–303. <https://doi.org/10.1080/17461391.2011.643923>.
94. Beck KL, Thomson JS, Swift RJ, Von Hurst PR. Role of nutrition in performance enhancement and postexercise recovery. *Open Access J Sports Med*. 2015;6:259–67. <https://doi.org/10.2147/OAJSM.S33605>.
95. O'Brien L, Collins K, Amirabdollahi F. Exploring sports nutrition knowledge in elite Gaelic footballers. *Nutrients*. 2021;13(4):1081. <https://doi.org/10.3390/nu13041081>.
96. Cassidy C, Collins K, Shortall M. The precompetition macronutrient intake of elite Gaelic football players. *Int J Sport Nutr Exerc Metab*. 2018;28(6):574–579. <https://doi.org/10.1123/ijsnem.2017-0292>.
97. Ó Catháin C, Fleming J, Renard M, Kelly D. Dietary intake of Gaelic football players during game preparation and recovery. *Sports*. 2020;8(5):62. <https://doi.org/10.3390/sports8050062>.
98. Martin L, Lambeth A, Scott D. Nutritional practices of national female soccer players: analysis and recommendations. *J Sports Sci Med*. 2006;5(1):130–7.
99. Condo D, Lohman R, Kelly M, Carr A. Nutritional Intake, sports nutrition knowledge and energy availability in female Australian rules football players. *Nutrients*. 2019;11(5):971. <https://doi.org/10.3390/nu11050971>.

100. Jenner SL, Devlin BL, Forsyth AK, Belski R. Dietary intakes of professional Australian football league women's (AFLW) athletes during a preseason training week. *J Sci Med Sport*. 2019;22(11):1266–71.
101. Williams C, Rollo I. Carbohydrate nutrition and team sport performance. *Sports Med*. 2015;45(1):S13–22.
102. Nattiv A, Loucks A, Manore M, Sanborn C, Sundgot-Borgen J, Warren M. American College of Sports Medicine position stand: the female athlete triad. *Med Sci Sports Exerc*. 2007;39(10):1867–82.
103. Mountjoy MJ, Sundgot-Borgen L, Burke S, Carter N, Constantini C, Lebrun N, Meyer R, Sherman K, Steffen KR, Budgett, Ljungqvist, A. The IOC consensus statement: Beyond the Female Athlete Triad-Relative Energy Deficiency in Sport (RED-S). *Br J Sports Med*. 2014;48(7):491–497.
104. Logue D, Madigan S, Heinen MM, Corish C. Screening for risk of low energy availability in athletic and recreationally active females in Ireland. *Eur J Sports Sci*. 2018;19(10):1–11. <https://doi.org/10.1080/17461391.2018.1526973>.
105. O'Brien L, Collins K, Doran D, Khaiyat O, Amirabdollahian F. Dietary intake and energy expenditure assessed during a pre-season period in elite Gaelic football players. *Sports (Basel)*. 2019;7(3):3. <https://doi.org/10.3390/sports7030062>.
106. Thomas DT, Erdman KA, Burke LM. American college of sports medicine joint position statement. nutrition and athletic performance. *Med Sci Sports Exerc*. 2016;48(3):543–568.
107. Russell M, Pennock A. Dietary analysis of young professional soccer players for 1 week during the competitive season. *J Strength Cond Res*. 2011;25(7):1816–23.
108. Cialdella-Kam L, Kulpins D, Manore MM. Vegetarian, gluten-free, and energy restricted diets in female athletes. *Sports* 2016;21(4):50. <https://doi.org/10.3390/sports4040050>.
109. Manore MM. Dietary recommendations and athletic menstrual dysfunction. *Sports Med*. 2002;32(14):887–901.
110. Manore MM. Weight management for athletes and active individuals: a brief review. *Sports Med*. 2015;45(1):83–92.
111. Ahmadi A, Enayatizadeh N, Akbarzadeh M, Asadi S, Tabatabaee SH. Iron status in female athletes participating in team ball-sports. *Pak J Biol Sci*. 2010;13(2):93–6.
112. Magee PJ, Gallagher AM, McCormack JM. High prevalence of dehydration and inadequate nutritional knowledge among university and club level athletes. *Int J Sport Nutr Exerc Metab*. 2017;27(2):158–68.
113. Todd J, Madigan S, Pourshahidi K, McSorley E, Laird E, Healy M, Magee P. Vitamin D status and supplementation practices in elite Irish athletes: an update from 2010/2011. *Nutrients*. 2016;8(8):485. <https://doi.org/10.3390/nu8080485>.
114. Gibbons E, Monagle, G Mc. Dietary intakes of calcium, iron and vitamin D in club and county female Gaelic footballers. Proceedings of the Nutrition Society. 2018; Volume 77, Issue OCE3: Irish Section Meeting, 20–22.
115. Aguilar-Navarro M, Baltazar-Martins G, Brito de Souza D, Muñoz-Guerra J, Del Mar Plata M, Del Coso J. Gender differences in prevalence and patterns of dietary supplement use in elite athletes. *Res Q Exerc Sport*. 2021;92(4):659–668.
116. Sawka MN, Burke LM, Eichner ER, Maughan RJ, Montain SJ, Stachenfeld NS. Exercise and fluid replacement. *Med Sci Sports Exerc*. 2007;39(2):377–90.
117. Ganio MS, Armstrong LE, Casa DJ, McDermott BP, Lee EC, Yamamoto LM, Marzano S, Lopez RM, Jimenez L, Le Bellego L, Chevillotte E, Lieberman HR. Mild dehydration impairs cognitive performance and mood of men. *Br J Nutr*. 2011;106(10):1535–43.
118. Maughan RJ, Shirreffs SM. Development of individual hydration strategies for athletes. *Int J Sports Nutr Exerc Metab*. 2008;18:457–72.
119. McCrink CM, McSorley EM, Grant K, McNeilly AM, Magee PJ. An investigation of dietary intake, nutrition knowledge and hydration status of Gaelic Football players. *Eur J Nutr*. 2021;60(3):1465–73.
120. Newell M, Newell J, Grant S. Fluid and electrolyte balance in elite Gaelic football players. *Ir Med J*. 2008;101(8):236–9.
121. Reilly T. The menstrual cycle and human performance: An overview. *Biol Rhy Res*. 2000;31(1):29–40.
122. Hewett TE, Myer GD, Ford KR. Anterior cruciate ligament injuries in female athletes: part 1, mechanisms and risk factors. *Am J Sports Med*. 2006;34(2):299–311.
123. Möller-Nielsen J, Hammar M. Women's soccer injuries in relation to the menstrual cycle and oral contraceptive use. *Med Sci Sports Exerc*. 1989;21(2):126–9.
124. Martin D, Timmins K, Cowie C, Alty J, Mehta R, Tang A, Varley I. Injury incidence across the menstrual cycle in international footballers. *Front Sports Act Living*. 2021;3:616999. <https://doi.org/10.3389/fspor.2021.616999>.
125. McNulty KL, Elliott-Sale KJ, Dolan E, Swinton PA, Ansdell P, Goodall S, Thomas K, Hicks KM. The effects of menstrual cycle phase on exercise performance in eumenorrheic women: a systematic review and meta-analysis. *Sports Med*. 2020;50(10):1813–27.
126. Elliott-Sale KJ, Minahan CL, de Jonge XAKJ, Ackerman KE, Sipilä S, Constantini NW, Lebrun CM, Hackney AC. Methodological considerations for studies in sport and exercise science with women as participants: a working guide for standards of practice for research on women. *Sports Med*. 2021;51(5):843–61.
127. Martin D, Sale C, Cooper SB, Elliott-Sale KJ. Period prevalence and perceived side effects of hormonal contraceptive use and the menstrual cycle in elite athletes. *Int J Sports Physiol Perform*. 2018;13(7):926–32.
128. Sims ST, Heather AK. Myths and Methodologies: Reducing scientific design ambiguity in studies comparing sexes and/or menstrual cycle phases. *Exp Physiol*. 2018;103:1309–17.
129. Elliott-Sale KJ, McNulty KL, Ansdell P, Goodall S, Hicks KM, Thomas K, Swinton PA, Dolan E. The effects of oral contraceptives on exercise performance in women: a systematic review and meta-analysis. *Sports Med*. 2020;50(10):1785–812.

Publisher's Note

Springer Nature remains neutral with regard to jurisdictional claims in published maps and institutional affiliations.

Ready to submit your research? Choose BMC and benefit from:

- fast, convenient online submission
- thorough peer review by experienced researchers in your field
- rapid publication on acceptance
- support for research data, including large and complex data types
- gold Open Access which fosters wider collaboration and increased citations
- maximum visibility for your research: over 100M website views per year

At BMC, research is always in progress.

Learn more biomedcentral.com/submissions

